Neuroscience Letters, 110 (1990) 356–360
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abcar, A., see Whitson, J.S., 319
Agid, Y., see Cervera, P., 210
Alkon, D.L., see Lederhendler, I.I., 28
Amann, R., see Maggi, C.A., 267
Arens, J., see Stabel, J., 60
Astolfi, M., see Maggi, C.A., 267
Avila, J., see Diaz-Nido, J., 221

Bakshi, R. and Faden, A.I., Blockade of the glycine modulatory site of NMDA receptors modifies dynorphin-induced behavioral effects, 113
Bauer, R., see Brüning, G., 6
Baumgarten, H.G., see Brüning, G., 6
Benistant, C., Rey, C. and Fonlupt, P., Increase of GABA-stimulated diazepam binding after lipid methylation in membrane preparations from rat brain, 137
Berkley, K.J., see Robbins, A., 82
Boczek-Funcke, A., Häbler, H.-J., Jänig, W. and Michaelis, M. Changes of activity in lumbar preganglionic neurones during slow urinary bladder filling in the cat, 97
Bornstein, J.C., see Hendriks, R., 286
Bourguignon, J.J., see Hechler, V., 204
Bragdon, A.C., see Mott, D.D., 131
Brüning, G., Bauer, R. and Baumgarten, H.G., Postnatal development of [^{3}H]flunitrazepam and [^{3}H]strychnine binding sites in rat spinal cord localized by quantitative autoradiography, 6
Burgess, G.M., see Dray, A., 52
Buzzell, G.R., see Mendendez-Pelaez, A., 314

Camp, D.M., see Robinson, T.E., 193
Carlson, M.D. and Ueda, T., Accumulated glutamate levels in the synaptic vesicle are not maintained in the absence of active transport, 325
Caudle, R.M., see Wimpey, T.L., 349
Caulfield, M.P., see Grove, E.A., 162
Cervera, P., Duyckaerts, C., Ruberg, M., Hirsch, E., Ransmayr, G., Hauw, J.-J. and Agid, Y., Tyrosine hydroxylase-like immunoreactivity in senile plaques is not related to the density of tyrosine hydroxylase-positive fibers in patients with Alzheimer's disease, 210
Chan, E.C., see Thomson, M., 343
Chavkin, C., see Wimpey, T.L., 349
Chen, L. and Huang, L.-Y.M., Ionic currents in retrogradely labeled trigeminothalamic neurons in slices of rat medulla, 66
Chihara, K., see Tsuruo, Y., 261
Claro, E., see Wallace, M.A., 155
Claustrat, B., see Vitte, P.A., 1
Collin, C., see Lederhendler, I.I., 28
Correa, F.M.A., Guilhaume, S.S. and Saavedra, J.M., Autoradiography of angiotensin-converting enzyme in fixed and unfixed rat brain using the specific enzyme inhibitor [^{125}I]351A or a polyclonal antibody and [^{125}I]staphylococcal protein A, 244
Cotman, C.W., see Whitson, J.S., 319

Daikoku, S., see Tsuruo, Y., 261
Davies, J., see Thomson, M., 343
De Barry, J., see Garcia-Ladona, F.J., 256
De Jong, W., see Lukovic, L., 46
Del Fiacco, M., Quartu, M., Floris, A. and Diaz, G., Substance P-like immunoreactivity in the human trigeminal ganglion, 16
Diaz, G., see Del Fiacco, M., 16
Diaz-Nido, J. and Avila, J., Aluminum induces the in vitro aggregation of bovine brain cytoskeletal proteins, 221
Dlampney, R.A.L., see McAllen, R.M., 91
Donnerer, J., see Maggi, C.A., 267
Dray, A., Forbes, C.A. and Burgess, G.M., Ruthenium red blocks the capsaicin-induced increase in intracellular calcium and activation of membrane currents in sensory neurones as well as the activation of peripheral nociceptors in vitro, 52
Dunwiddie, T.V., see Su, M.-T., 186
Duyckaerts, C., see Cervera, P., 210

Eaton, S.A. and Salt, T.E., Thalamic NMDA receptors and nociceptive sensory synaptic transmission, 297
Evans, F.J., see Grove, E.A., 162
Eysel, U.Th., see Vandenbussche, E., 249

Faden, A.I., see Bakshi, R., 113
Faden, A.I., see Graham, S.H., 124
Falconer, J., see Thomson, M., 343
Fearn, S., see Heafield, M.T., 216
Ferre, S., see Herrera-Marschitz, M., 172
Floris, A., see Del Fiacco, M., 16
Foehring, R.C. and Wyler, A.R., Two patterns of firing in human neocortical neurons, 279
Fonlupt, P., see Benistant, C., 137
Forbes, C.A., see Dray, A., 52

0304-3940/90/$ 03.50 © 1990 Elsevier Scientific Publishers Ireland Ltd.

Fujii, T., Hayashi, M., Toita, K.-i. and Yamasaki, Y., Effects of coenzymes Q_2 and Q_{10} on the field potential of guinea pig olfactory cortex slices maintained in hypoxia, 40
Furness, J.B., see Hendriks, R., 286

Garcia-Ladona, F.J., Palacios, J.M., De Barry, J. and Gombos, G., Developmentally regulated changes of glutamate binding sites in mouse deep cerebellar nuclei, 256
Ge, L.-J., see Zadina, J.E., 143
Geraghty, S., see Thomson, M., 343
Gerhardt, G.A., see Su, M.-T., 186
Ghodsi-Hovsepian, S., Messer, W.S. Jr. and Hoss, W., Subtype specificity of the muscarinic receptor-stimulated GTPase response in the rat cortex, 148
Glabe, C.G., see Whitson, J.S., 319
Goiny, M., see Herrera-Marschitz, M., 172
Gombos, G., see Garcia-Ladona, F.J., 256
Gonzalez-Mora, J.L., see Mas, M., 303
Gottmann, K. and Lux, H.D., Low- and high-voltage-activated Ca^{2+} conductances in electrically excitable growth cones of chick dorsal root ganglion neurons, 34
Graham, S.H., Shiraishi, K., Panter, S.S., Simon, R.P. and Faden, A.I., Changes in extracellular amino acid neurotransmitters produced by focal cerebral ischemia, 124
Grove, E.A., Caulfield, M.P. and Evans, F.J., Inhibition of protein kinase C prevents phorbol ester- but not muscarine-induced depolarizations in the rat superior cervical ganglion, 162
Guadalupe, T., see Mas, M., 303
Guilhaume, S.S., see Correa, F.M.A., 244

Häbler, H.-J., see Boczek-Funcke, A., 97
Hakansson, L., see Herrera-Marschitz, M., 172
Hanada, M., see Tamiya, R., 11
Harthe, C., see Vitte, P.A., 1
Hashimoto, S., Kawamura, J., Segawa, Y., Suenaga, T. and Nakamura, M., Bifurcation of P9 far-field potentials induced by changes in the shoulder position, 102
Hattori, Y., see Moriwaki, A., 72
Hauw, J.-J., see Cervera, P., 210
Hayashi, M., see Fujii, T., 40
Heafield, M.T., Fearn, S., Steventon, G.B., Waring, R.H., Williams, A.C. and Sturman, S.G., Plasma cysteine and sulphate levels in patients with motor neurone, Parkinson's and Alzheimer's disease, 216
Hechler, V., Schmitt, M., Bourguignon, J.J. and Miatre, M., Trans-γ-hydroxycrotonic acid binding sites in brain: evidence for a subpopulation of γ-hydroxybutyrate sites, 204
Heinemann, U., see Stabel, J., 60
Hendriks, R., Bornstein, J.C. and Furness, J.B., An electrophysiological study of the projections of putative sensory neurons within the myenteric plexus of the guinea pig ileum, 286
Henke, P.G., see Ray, A., 331
Herbst, W.M., see Kummer, W., 239
Herrera-Marschitz, M., Goiny, M., Utsumi, H., Ferre, S., Hakansson, L., Nordberg, A. and Ungerstedt, U., Effect of unilateral nucleus basalis lesion on cortical and striatal acetylcholine and dopamine release monitored in vivo with microdialysis, 172
Heym, C., see Kummer, W., 239
Hirsch, E., see Cervera, P., 210
Hisano, S., see Tsuruo, Y., 261
Höllt, V., see Moneta, M.E., 273
Hori, Y., see Moriwaki, A., 72
Hoss, W., see Ghodsi-Hovsepian, S., 148
Hotta, H., see Robbins, A., 82
Huang, L.-Y.M., see Chen, L., 66
Humlová, M., see Illnerová, H., 77
Hurtado, H., see Knoops, B., 180

Illnerová, H. and Humlová, M., The rat pineal *N*-acetyltransferase rhythm persists after a five-hour, but disappears temporarily after a seven-hour advance of the light–dark cycle: a six-hour shift may be a turning point, 77
Inagaki, S., see Tamiya, R., 11

Jensen, M.S., see Nathan, T., 309
Jänig, W., see Boczek-Funcke, A., 97

Kagotani, Y., see Tsuruo, Y., 261
Kamiya, H.-o., see Saito, R., 337
Kantalrjian, A., see Seidel, A., 227
Kastin, A.J., see Zadina, J.E., 143
Kawae, N., see Ohara-Imaizumi, M., 167
Kawai, Y., see Tamiya, R., 11
Kawamura, J., see Hashimoto, S., 102
Kawano, H., see Tsuruo, Y., 261
Kikuchi, H., see Nozaki, K., 22
Knoops, B., Hurtado, H. and Van den Bosch de Aguilar, Ph., Effects of dibutyryl cyclic adenosine monophosphate, ACTH(4–10) and gangliosides on nerve fiber regeneration after rat sciatic nerve tubulization, 180

Konishi, H., see Saito, R., 337
Kumakura, K., see Ohara-Imaizumi, M., 167
Kummer, W., Herbst, W.M. and Heym, C., Vasoactive intestinal polypeptide receptor-like immunoreactivity in human sweat glands, 239

Lambert, J.D.C., see Nathan, T., 309
Lambert, J.D.C., see Stabel, J., 60
Lambert, J.J., see Peters, J.A., 107
Lederhendler, I.I., Collin, C. and Alkon, D.L., Sequential changes of potassium currents in *Hermissenda* Type B photoreceptor during early stages of classical conditioning, 28
Levy, D.I., Sucher, N.J. and Lipton, S.A., Redox modulation of NMDA receptor-mediated toxicity in mammalian central neurons, 291
Lewis, D.V., see Mott, D.D., 131
Lipton, P., see Raley, K.M., 118
Lipton, S.A., see Levy, D.I., 291
Louilot, A., see Mas, M., 303
Lukovic, L. and De Jong, W., Analysis of the cardiovascular changes evoked by microinjection of NaCl into the nucleus tractus solitarii of rats: evidence for a Na^+/Ca^{2+} relationship, 46
Lux, H.D., see Gottmann, K., 34

Maggi, C.A., Astolfi, M., Donnerer, J. and Amann, R., Which mechanisms account for the sensory neuron blocking action of capsaicin of primary afferents in the rat urinary bladder?, 267
Malone, H.M., see Peters, J.A., 107
Mas, M., Gonzalez-Mora, J.L., Louilot, A., Solé, C. and Guadalupe, T., Increased dopamine release in the nucleus accumbens of copulating male rats as evidenced by in vivo voltammetry, 303
McAllen, R.M. and Dlampney, R.A.L., Vasomotor neurons in the rostral ventrolateral medulla are organized topographically with respect to type of vascular bed but not body region, 91
Mendendez-Pelaez, A., Buzzell, G.R., Nonaka, K.O. and Reiter, R.J. In vivo administration of isoproterenol or forskolin during the light phase induces increases in the melatonin content of the Syrian hamster pineal gland without a rise in *N*-acetyltransferase activity, 314
Messer, W.S. Jr., see Ghodsi-Hovsepian, S., 148
Miatre, M., see Hechler, V., 204
Michaelis, M., see Boczek-Funcke, A., 97
Mizuno, N., see Nozaki, K., 22
Moneta, M.E. and Höllt, V., Perforant path kindling induces differential alterations in the mRNA levels coding for prodynorphin and proenkephalin in the rat hippocampus, 273
Moriwaki, A., Hattori, Y., Nishida, N. and Hori, Y., Electrocorticographic characterization of chronic iron-induced epilepsy in rats, 72
Mott, D.D., Bragdon, A.C. and Lewis, D.V., Phaclofen antagonizes post-tetanic disinhibition in the rat dentate gyrus, 131
Mynlieff, M., see Su, M.-T., 186

Nakamura, M., see Hashimoto, S., 102
Nathan, T., Jensen, M.S. and Lambert, J.D.C., The slow inhibitory postsynaptic potential in rat hippocampal CA1 neurones is blocked by intracellular injection of QX-314, 309
Nishida, N., see Moriwaki, A., 72
Nonaka, K.O., see Mendendez-Pelaez, A., 314
Nonaka, S., see Saito, R., 337
Nordberg, A., see Herrera-Marschitz, M., 172
Nozaki, K., Okamoto, S., Uemura, Y., Kikuchi, H. and Mizuno, N., Erythrocytes suppress calcitonin gene-related peptide- and vasoactive intestinal polypeptide-like immunoreactivities in cerebrovascular nerve fibers after subarachnoid hemorrhage, 22

Ohara-Imaizumi, M., Takeda, K., Kawae, N. and Kumakura, K., Effects of pertussis toxin on the affinity of exocytosis for Ca^{2+} in bovine adrenal chromaffin cells, 167
Okamoto, S., see Nozaki, K., 22
Orban, G.A., see Vandenbussche, E., 249

Palacios, J.M., see Garcia-Ladona, F.J., 256
Panter, S.S., see Graham, S.H., 124
Paschen, W., Röhn, G. and Schmidt-Kastner, R., Transglutaminase activity in reversible cerebral ischemia in the rat, 232
Paulson, P.E., see Robinson, T.E., 193
Peters, J.A., Malone, H.M. and Lambert, J.J., Antagonism of 5-HT_3 receptor mediated currents in murine N1E-115 neuroblastoma cells by (+)-tubocurarine, 107
Pevet, P., see Vitte, P.A., 1
Pubols, B.H., Jr., Slowly adapting Type I mechanoreceptor discharge as a function of dynamic force versus dynamic displacement of glabrous skin of raccoon and squirrel monkey hand, 86

Quartu, M., see Del Fiacco, M., 16

Raley, K.M. and Lipton, P., NMDA receptor activation accelerates ischemic energy depletion in the hippocampal slice and the demonstration of a threshold for ischemic damage to protein synthesis, 118
Ransmayr, G., see Cervera, P., 210
Ray, A., Henke, P.G. and Sullivan, R.M., Noradrenergic mechanisms in the central amygdalar nucleus and gastric stress ulcer formation in rats, 331
Reiter, R.J., see Mendendez-Pelaez, A., 314
Rey, C., see Benistant, C., 137
Robbins, A., Sato, Y., Hotta, H. and Berkley, K.J., Responses of hypogastric nerve afferent fibers to uterine distension in estrous or metestrous rats, 82
Robinson, T.E., Yew, J., Paulson, P.E. and Camp, D.M., The long-term effects of neurotoxic doses of methamphetamine on the extracellular concentration of dopamine measured with microdialysis in striatum, 193
Röhn, G., see Paschen, W., 232
Ruberg, M., see Cervera, P., 210

Saavedra, J.M., see Correa, F.M.A., 244
Saito, R., Konishi, H., Takano, Y., Nonaka, S., Sakaguchi, K., Shimohigashi, Y. and Kamiya, H.-o., Characterization of tachykinin receptors in endothelial cells of porcine artery, 337
Sakaguchi, K., see Saito, R., 337
Salt, T.E., see Eaton, S.A., 297
Sato, Y., see Robbins, A., 82
Schmidt-Kastner, R., see Paschen, W., 232
Schmitt, M., see Hechler, V., 204
Segawa, Y., see Hashimoto, S., 102
Seidel, A., Kantalrjian, A. and Vollrath, L., A possible role for cyclic guanosine monophosphate in the rat pineal gland, 227
Shimohigashi, Y., see Saito, R., 337
Shintani, E., see Whitson, J.S., 319
Shiraishi, K., see Graham, S.H., 124
Simon, R.P., see Graham, S.H., 124
Smith, R., see Thomson, M., 343
Solé, C., see Mas, M., 303
Stabel, J., Arens, J., Lambert, J.D.C. and Heinemann, U., Effects of lowering $[Na^+]_o$ and $[K^+]_o$ and of ouabain on quisqualate-induced ionic changes in area CA1 of rat hippocampal slices, 60
Steventon, G.B., see Heafield, M.T., 216
Sturman, S.G., see Heafield, M.T., 216
Su, M.-T., Dunwiddie, T.V., Mynlieff, M. and Gerhardt, G.A., Electrochemical characterization of stimulated norepinephrine overflow in locus coeruleus-hippocampus double brain grafts grown in oculo, 186
Sucher, N.J., see Levy, D.I., 291
Suenaga, T., see Hashimoto, S., 102
Sullivan, R.M., see Ray, A., 331

Takagi, H., see Tamiya, R., 11
Takano, Y., see Saito, R., 337
Takeda, K., see Ohara-Imaizumi, M., 167
Tamiya, R., Hanada, M., Kawai, Y., Inagaki, S. and Takagi, H., Substance P afferents have synaptic contacts with dopaminergic neurons in the ventral tegmental area of the rat, 11
Thomson, M., Chan, E.C., Davies, J., Falconer, J., Geraghty, S. and Smith, R., Intracellular mechanisms governing the acute phase of β-endorphin secretion from the corticotrope in vitro, 343
Toita, K.-i., see Fujii, T., 40
Tsuruo, Y., Kawano, H., Kagotani, Y., Hisano, S., Daikoku, S., Chihara, K., Zhang, T. and Yanaihara, N., Morphological evidence for neuronal regulation of luteinizing hormone-releasing hormone-containing neurons by neuropeptide Y in the rat septo-preoptic area, 261

Ueda, T., see Carlson, M.D., 325
Uemura, Y., see Nozaki, K., 22
Ungerstedt, U., see Herrera-Marschitz, M., 172
Utsumi, H., see Herrera-Marschitz, M., 172

Van den Bosch de Aguilar, Ph., see Knoops, B., 180
Vandenbussche, E., Eysel, U.Th. and Orban, G.A., Influence of retinal lesions on grating acuity of the cat, 249
Vanecek, J. and Vollrath, L., Melatonin modulates diacylglycerol and arachidonic acid metabolism in the anterior pituitary of immature rats, 199
Vitte, P.A., Harthe, C., Pevet, P. and Claustrat, B., Brain autoradiographic study in the golden hamster after intracarotid injection of [^{14}H]melatonin, 1
Vollrath, L., see Seidel, A., 227
Vollrath, L., see Vanecek, J., 199

Wallace, M.A. and Claro, E., A novel role for dopamine: inhibition of muscarinic cholinergic-stimulated phosphoinositide hydrolysis in rat brain cortical membranes, 155

Waring, R.H., see Heafield, M.T., 216
Whitson, J.S., Glabe, C.G., Shintani, E., Abcar, A. and Cotman, C.W., β-Amyloid protein promotes neuritic branching in hippocampal cultures, 319
Williams, A.C., see Heafield, M.T., 216
Wimpey, T.L., Caudle, R.M. and Chavkin, C., Chronic morphine exposure blocks opioid effects on both the early and late inhibitory postsynaptic potential in hippocampal CA1 pyramidal cells, 349
Wyler, A.R., see Foehring, R.C., 279

Yamasaki, Y., see Fujii, T., 40
Yanaihara, N., see Tsuruo, Y., 261
Yew, J., see Robinson, T.E., 193

Zadina, J.E., Kastin, A.J. and Ge, L.-J., TYR-MIF-1 binding in brain is not altered by ligands selective for the $GABA_A$/benzodiazepine receptor, 143
Zhang, T., see Tsuruo, Y., 261

CONTENTS *(continued from back cover)*

Characterization of tachykinin receptors in endothelial cells of porcine artery
R. Saito, H. Konishi, Y. Takano, S. Nonaka, K. Sakaguchi, Y. Shimohigashi and H.-o. Kamiya (Japan) 337
Intracellular mechanisms governing the acute phase of β-endorphin secretion from the corticotrope in vitro
M. Thomson, E.C. Chan, J. Davies, J. Falconer, S. Geraghty and R. Smith 343
Chronic morphine exposure blocks opioid effects on both the early and late inhibitory postsynaptic potential in hippocampal CA1 pyramidal cells
T.L. Wimpey, R.M. Caudle and C. Chavkin (U.S.A.) 349

Author Index 356